

**Figure 1B: siNA Hybridization Assay**





Figure 1C: Principle of siNA detection/quantitation Assays



[C] - [A] = [Duplex] based upon analysis of sense strand  
[D] - [B] = [Duplex] based upon analysis of antisense strand  
[E] = [Duplex] based upon direct analysis of duplex

Figure 2A: siNA Stab 7 Sense Strand Standard Curve






Figure 2B: siNA Stab 8 Antisense Strand Standard Curve






**Figure 2C: siNA Stab 7 Cholesterol Conjugate  
Sense Strand Standard Curve**




**Figure 2D: siNA Stab 7 Trigalactose Cholesterol  
Conjugate Antisense Strand Standard Curve**



8/27


Figure 3A: siNA Stab 7 Single Stranded  
Quality Control Sample






9/27

**Figure 3B: siNA Stab 8 Single Stranded  
Quality Control Sample**




10/27

**Figure 3C: siNA Stab 7 Duplex  
Quality Control Sample**




**Figure 4A: Detection of potential  
siNA Stab 7 Metabolites via Hybridization Assay**






12/27

**Figure 4B: Detection of potential  
siNA Stab 8 Metabolites via Hybridization Assay**






**Figure 5A: Effect of Hepatocyte lysate on detection of single stranded Stab 7 cholesterol conjugate siNA sequence**




**Figure 5B: Effect of Hepatocyte lysate on detection of single stranded Stab 8 siNA sequence**






**Figure 5C: Effect of Hepatocyte lysate on detection of Stab 7 cholesterol conjugate duplex siNA sequence**




**Figure 6A: Effect of monkey plasma on detection of single stranded Stab 7 cholesterol conjugate siNA sequence**






**Figure 6B: Effect of monkey plasma on detection of single stranded Stab 8 siNA sequence**




**Figure 6C: Effect of monkey plasma on detection of Stab 7 cholesterol conjugate duplex siNA sequence**




19/27

**Figure 7: Concentration of siNA duplex and antisense  
In Hepatocytes**




Given - 1 gram liver =  $1 \text{ mL} = 10^{10}$  cells      [siRNA ~  $\mu\text{M}$ ]

**Figure 8: Removal of Competitive binding sequence  
In duplex assay**






**Figure 9: Application of Hybridization Assay to siNA molecules having identical sequence with differing chemical modifications**





22/27

Figure 10



23/27

Figure 11




24/27

Figure 12


... instruction of -Q1: 8.643 to 11.023 min from Sample 1 (Sample001) of duplex050903002.wiff

Max. 1.0 cps.



Mass reconstruction of -Q1: 9.520 to 10.981 min from Sample 2 (Sample002) of duplex051603001.wiff

Max. 1.0 cps.





25/27

Figure 13







Figure 14



27/27

**Figure 15**

In this design, binding of the target siRNA (in step 1) prevents binding of a secondary detection probe (in step 2). Therefore, signal is inversely proportional to analyte concentration.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**